

Final RASML entries for mometasone, with changes to previously proposed entries included as tracked changes:

Column 1 Substance(s)	Column 2 Conditions	Column 3 Required statement(s)
Mometasone (Entry 1 of 2)	In preparations for dermal use	<ul style="list-style-type: none"> • Do not use: <ul style="list-style-type: none"> - in or around the eyes - for acne - on broken skin or skin infections (eg. cold sores, shingles, chicken pox, thrush, tinea, ringworm) or ulcerous skin - if you are sensitive to any corticosteroids • Unless a doctor has told you to, do not use: <ul style="list-style-type: none"> - on the face, large areas of the skin, or areas that rub together - under a bandage, dressing, nappies or plastic pants - for more than 7 days - if you are pregnant, likely to become pregnant or breastfeeding - for children under 12 years old • If irritation occurs, discontinue use and seek medical advice • Seek medical advice if there is no improvement in symptoms after using for 1 week
Mometasone (Entry 2 of 2)	In nasal spray preparations	<ul style="list-style-type: none"> • Do not exceed the maximum stated dose. • Do not use for more than 6 months without the advice of your doctor or pharmacist • See your doctor or pharmacist before using this product if: <ul style="list-style-type: none"> - you have a nasal or sinus infection - you have recently had an injury or surgery to your nose - you have ulceration in your nose - you are pregnant or likely to become pregnant or are breastfeeding • See your doctor or pharmacist if: <ul style="list-style-type: none"> - symptoms are not relieved within 7 days - your nose bleeds - you develop signs or symptoms of a nasal or sinus infection such as fever, pain or swelling, or discoloured nasal discharge - you have eye pain or visual disturbances