

Final RASML entries for OTC triptans with changes to previously proposed entries included as tracked changes:

Column 1 Substance(s)	Column 2 Conditions	Column 3 Required statement(s)
<p>Sumatriptan Rizatriptan <u>Zolmitriptan</u>* (Entry 1 of 1)</p> <p>* These will have individual but identical entries</p>	<p>In divided oral preparations for relief of migraine</p>	<ul style="list-style-type: none"> • Do not use: <ul style="list-style-type: none"> - without a previous diagnosis of migraine by a doctor - if you have previously had a heart attack or stroke or if you have heart disease, peripheral vascular disease (a blood vessel disorder) or high blood pressure or have previously had a heart attack or stroke. — if you have taken other migraine medications within 24 hours — if you have taken monoamine oxidase inhibitors (MAOIs; medicines used to treat depression) in the last 2 weeks — if you have liver or kidney problems, unless advised by your doctor • Consult your doctor or pharmacist before use: <ul style="list-style-type: none"> - if you think you may be at risk of heart disease (including if you are post-menopausal or a male over the age of 40) - if you have <u>liver or kidney problems or</u> any other medical conditions - <u>if you are taking other medicines for migraine</u> - if you are taking <u>or have taken in the last two weeks, medicine</u> medication for depression <u>or anxiety</u> - if you are pregnant or breastfeeding • Speak to a doctor or pharmacist if your migraines <u>attacks</u> become more frequent. Overuse of this medicine may worsen your condition. • This medicine may cause drowsiness. If affected, do not drive a vehicle or operate machinery.
<p>Eletriptan (Entry 1 of 1)</p>	<p>In divided oral preparations for relief of migraine</p>	<ul style="list-style-type: none"> • Do not use: <ul style="list-style-type: none"> - without a previous diagnosis of migraine by a doctor - if you have previously had a heart attack or stroke or if you have heart disease, peripheral vascular disease (a blood vessel disorder) or high blood pressure or have previously had a heart attack or stroke. — if you have taken other migraine medications within 24 hours — if you have liver or kidney problems, unless advised by your doctor • Consult your doctor or pharmacist before use: <ul style="list-style-type: none"> - if you think you may be at risk of heart disease (including if you are post-menopausal or a male over the age of 40) - if you have <u>liver or kidney problems or</u> any other medical conditions - if you are taking <u>any other medicines, including for migraine,</u> medication for depression <u>or anxiety</u> - if you are pregnant or breastfeeding • Speak to a doctor or pharmacist if your migraines <u>attacks</u> become more frequent. Overuse of this medicine may worsen your condition. • This medicine may cause drowsiness. If affected, do not drive a vehicle or operate machinery.

Final RASML entries for OTC triptans (without tracked changes):

Column 1 Substance(s)	Column 2 Conditions	Column 3 Required statement(s)
<p>Sumatriptan Rizatriptan Zolmitriptan* (Entry 1 of 1)</p> <p>* These will have individual but identical entries</p>	<p>In divided oral preparations for relief of migraine</p>	<ul style="list-style-type: none"> • Do not use: <ul style="list-style-type: none"> - without a previous diagnosis of migraine by a doctor - if you have heart disease, peripheral vascular disease (a blood vessel disorder) or high blood pressure or have previously had a heart attack or stroke. • Consult your doctor or pharmacist before use: <ul style="list-style-type: none"> - if you think you may be at risk of heart disease (including if you are post-menopausal or a male over the age of 40) - if you have liver or kidney problems or any other medical conditions - if you are taking other medicines for migraine - if you are taking, or have taken in the last two weeks, medicine for depression or anxiety - if you are pregnant or breastfeeding • Speak to a doctor or pharmacist if your migraine attacks become more frequent. Overuse of this medicine may worsen your condition. • This medicine may cause drowsiness. If affected, do not drive a vehicle or operate machinery.
<p>Eletriptan (Entry 1 of 1)</p>	<p>In divided oral preparations for relief of migraine</p>	<ul style="list-style-type: none"> • Do not use: <ul style="list-style-type: none"> - without a previous diagnosis of migraine by a doctor - if you have heart disease, peripheral vascular disease (a blood vessel disorder) or high blood pressure or have previously had a heart attack or stroke. • Consult your doctor or pharmacist before use: <ul style="list-style-type: none"> - if you think you may be at risk of heart disease (including if you are post-menopausal or a male over the age of 40) - if you have liver or kidney problems or any other medical conditions - if you are taking any other medicines, including for migraine, depression or anxiety - if you are pregnant or breastfeeding • Speak to a doctor or pharmacist if your migraine attacks become more frequent. Overuse of this medicine may worsen your condition. • This medicine may cause drowsiness. If affected, do not drive a vehicle or operate machinery.